

WHAT IS CLAIMED IS:

1. A composition comprising an isolated or recombinant peptide comprising a subsequence of a Class II major histocompatibility molecule, wherein the peptide has the following properties,

5 (a) having a structure comprising $R_1 - R_2 - R_3 - R_4 - R_5 - R_6 - R_7 - R_8 - R_9 - R_{10} - R_{11} - R_{12} - R_{13} - R_{14} - R_{15} - R_{16}$,

wherein R_1 is Gln, Lys, or Arg; R_2 is Arg; R_3 and R_4 are members independently selected from the group consisting of all amino acids; R_5 is Ala, Glu, Asp, Val, Leu or Ile; R_6 and R_7 are members independently selected from the group consisting of
 10 all amino acids; R_8 is Thr; $R_9, R_{10}, R_{11}, R_{12}, R_{13}, R_{14}$, and R_{15} are members independently selected from the group consisting of all amino acids; and, R_{16} is Val;

(b) capable of generating an immune response to a non-Hodgkin's B cell lymphoma cell.

15 2. The composition of claim 1, wherein R_1 is Gln, Lys, or Arg; R_2 is Arg; R_3 is Arg; R_4 is selected from the group consisting of all amino acids; R_5 is Ala; R_6 and R_7 are members independently selected from the group consisting of all amino acids; R_8 is Thr; R_9 is selected from the group consisting of all amino acids; R_{10} is Cys; $R_{11}, R_{12}, R_{13}, R_{14}$, and R_{15} are members independently selected from the group consisting of all amino acids; and, R_{16} is Val.

20 3. The composition of claim 2, wherein R_1 is Gln, Lys, or Arg; R_2 is Arg; R_3 is Arg; R_4 is Ala; R_5 is Ala; R_6 is Val; R_7 is Asp; R_8 is Thr; R_9 is Tyr; R_{10} is Cys; R_{11} is Arg; R_{12} is His; R_{13} is Asn; R_{14} is Tyr; R_{15} is Gly, and R_{16} is Val.

25 4. The composition of claim 1, further comprising a pharmaceutically acceptable excipient.

5. The composition of claim 1, further comprising an adjuvant.

30 6. The composition of claim 1, wherein the non-Hodgkin's lymphoma cell is selected from the group consisting of a B-cell chronic lymphocytic leukemia/small lymphocytic lymphoma (B-CCL/SLL) cell, a lymphoplasmacytoid lymphoma (LPL) cell, a follicular

lymphoma (FL) cell, a mucosa-associated lymphoid tissue lymphoma (MALTL) cell, a splenic lymphoma with villous lymphocytes (SLVL) cell and a mantle cell lymphoma cell.

7. A method for detecting a nucleic acid in a biological sample, wherein the nucleic acid encodes a peptide capable of specifically binding to a Lym-1 antibody, the method comprising the following steps:
 - (a) contacting the sample with an oligonucleotide primer pair capable of amplifying a subsequence of an MHC nucleic acid, which subsequence encodes a polypeptide comprising a peptide of claim 1,
 - (b) amplifying the nucleic acid; and
 - (c) detecting the amplified nucleic acid.
8. The method of claim 7, wherein the MHC gene is HLA-DR 10.
9. The method of claim 7, wherein the subsequence encodes a peptide wherein R₁ is Gln, Lys, or Arg; R₂ is Arg; R₃ is Arg; R₄ is Ala; R₅ is Ala; R₆ is Val; R₇ is Asp; R₈ is Thr; R₉ is Tyr; R₁₀ is Cys; R₁₁ is Arg; R₁₂ is His; R₁₃ is Asn; R₁₄ is Tyr; R₁₅ is Gly, and R₁₆ is Val.
10. The method of claim 7, wherein the biological sample comprises a B cell.
11. The method of claim 10, wherein the B cell is a B lymphocytic non-Hodgkin's lymphoma cell.
12. The method of claim 11, wherein the non-Hodgkin's lymphoma cell is selected from the group consisting of a B-cell chronic lymphocytic leukemia/small lymphocytic lymphoma (B-CCL/SLL) cell, a lymphoplasmacytoid lymphoma (LPL) cell, a follicular lymphoma (FL) cell, a mucosa-associated lymphoid tissue lymphoma (MALTL) cell, a splenic lymphoma with villous lymphocytes (SLVL) cell and a mantle cell lymphoma cell.
13. The method of claim 7, wherein the biological sample is a body fluid sample or a biopsy sample.
14. The method of claim 13, wherein the body fluid sample is a blood sample.

15. A kit for detecting a nucleic acid in a biological sample, wherein the nucleic acid encodes a peptide capable of specifically binding to a Lym-1 antibody, comprising an oligonucleotide primer pair capable of amplifying a subsequence of an MHC gene or gene product, which subsequence encodes a polypeptide comprising a peptide of claim 1.

5

16. The kit of claim 15, wherein the MHC gene is HLA-DR 10.

10

17. The kit of claim 15, wherein R₁ is Gln, Lys, or Arg; R₂ is Arg; R₃ is Arg; R₄ is Ala; R₅ is Ala; R₆ is Val; R₇ is Asp; R₈ is Thr; R₉ is Tyr; R₁₀ is Cys; R₁₁ is Arg; R₁₂ is His; R₁₃ is Asn; R₁₄ is Tyr; R₁₅ is Gly, and R₁₆ is Val.

15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995

18. The kit of claim 15, further comprising an instructional material teaching a use of the kit, wherein the instructional material indicates that the kit is used for the detection of nucleic acid encoding a peptide reactive with a Lym-1 antibody and that the polypeptide is associated with non-Hodgkin's B cell lymphomas.

19. A method for detecting an antibody reactive with a non-Hodgkin's B cell lymphoma cell, comprising:

- (a) contacting a sample with a composition of claim 1 under immunologically reactive conditions, and
- (a) detecting whether an antibody has specifically bound to the composition.

20. The method of claim 19, wherein the sample is a biological sample.

21. The method of claim 19, wherein R₁ is Gln, Lys, or Arg; R₂ is Arg; R₃ is Arg; R₄ is Ala; R₅ is Ala; R₆ is Val; R₇ is Asp; R₈ is Thr; R₉ is Tyr; R₁₀ is Cys; R₁₁ is Arg; R₁₂ is His; R₁₃ is Asn; R₁₄ is Tyr; R₁₅ is Gly, and R₁₆ is Val.

22. The method of claim 19, wherein the antibody is generated by a recombinant nucleic acid library.

23. The method of claim 22, wherein the recombinant nucleic acid is a phage display library.

24. The method of claim 19, wherein the composition is fixed to a solid surface.

25. A method for generating an antibody reactive with a non-Hodgkin's B cell lymphoma cell, comprising administering an immunogenically effective amount of a composition of
5 claim 1 to a mammal. J

26. The method of claim 22, wherein R₁ is Gln, Lys, or Arg; R₂ is Arg; R₃ is Arg; R₄ is Ala; R₅ is Ala; R₆ is Val; R₇ is Asp; R₈ is Thr; R₉ is Tyr; R₁₀ is Cys; R₁₁ is Arg; R₁₂ is His; R₁₃ is Asn; R₁₄ is Tyr; R₁₅ is Gly, and R₁₆ is Val.

10

27. The method of claim 25, wherein the non-Hodgkin's lymphoma cell is selected from the group consisting of a B-cell chronic lymphocytic leukemia/small lymphocytic lymphoma (B-CCL/SLL) cell, a lymphoplasmacytoid lymphoma (LPL) cell, a follicular lymphoma (FL) cell, a mucosa-associated lymphoid tissue lymphoma (MALTL) cell, a splenic lymphoma with villous lymphocytes (SLVL) cell and a mantle cell lymphoma cell.

5

28. An immunogenic composition capable of eliciting an immunogenic response directed to a polypeptide epitope, wherein the epitope comprises an amino acid sequence having a structure comprising

20

R₁ - R₂ - R₃ - R₄ - R₅ - R₆ - R₇ - R₈ - R₉ - R₁₀ - R₁₁ - R₁₂ - R₁₃ - R₁₄ - R₁₅ - R₁₆,

wherein R₁ is Gln, Lys, or Arg; R₂ is Arg; R₃ and R₄ are members

independently selected from the group consisting of all amino acids; R₅ is Ala, Glu, Asp, Val, Leu or Ile; R₆ and R₇ are members independently selected from the group consisting of all amino acids; R₈ is Thr; R₉, R₁₀, R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅ are members independently
25 selected from the group consisting of all amino acids; and, R₁₆ is Val.

25

29. The immunogenic composition of claim 28, wherein R₁ is Gln, Lys, or Arg; R₂ is Arg; R₃ is Arg; R₄ is Ala; R₅ is Ala; R₆ is Val; R₇ is Asp; R₈ is Thr; R₉ is Tyr; R₁₀ is Cys; R₁₁ is Arg; R₁₂ is His; R₁₃ is Asn; R₁₄ is Tyr; R₁₅ is Gly, and R₁₆ is Val.

30

30. The immunogenic composition of claim 28, wherein the immunogenic response generates antibodies specific for the polypeptide epitope.

31. A method of inducing an immunogenic response directed to a polypeptide epitope, comprising administering an immunogenically effective amount of a composition comprising a polypeptide epitope to a mammal,

wherein the epitope comprises an amino acid sequence having a structure

comprising $R_1 - R_2 - R_3 - R_4 - R_5 - R_6 - R_7 - R_8 - R_9 - R_{10} - R_{11} - R_{12} - R_{13} - R_{14} - R_{15} - R_{16}$,

wherein R_1 is Gln, Lys, or Arg; R_2 is Arg; R_3 and R_4 are members

independently selected from the group consisting of all amino acids; R_5 is Ala, Glu, Asp,

Val, Leu or Ile; R_6 and R_7 are members independently selected from the group consisting of

all amino acids; R_8 is Thr; $R_9, R_{10}, R_{11}, R_{12}, R_{13}, R_{14}$, and R_{15} are members independently

selected from the group consisting of all amino acids; and, R_{16} is Val.

32. The method of claim 31, wherein R_1 is Gln, Lys, or Arg; R_2 is Arg; R_3 is Arg; R_4 is Ala; R_5 is Ala; R_6 is Val; R_7 is Asp; R_8 is Thr; R_9 is Tyr; R_{10} is Cys; R_{11} is Arg; R_{12} is His; R_{13} is Asn; R_{14} is Tyr; R_{15} is Gly, and R_{16} is Val.

33. The immunogenic composition of claim 31, wherein the immunogenic response generates antibodies specific for the polypeptide epitope.

34. The method of claim 31, wherein the mammal is a human, a mouse or a rabbit.